

SUPER SOLUTIONS OF THE DYNAMICAL YANG-BAXTER EQUATION

GIZEM KARAALI

ABSTRACT. A super dynamical r -matrix r satisfies the *zero weight condition* if:

$$[h \otimes 1 + 1 \otimes h, r(\lambda)] = 0 \text{ for all } h \in \mathfrak{h}, \lambda \in \mathfrak{h}^*.$$

In this paper we classify super dynamical r -matrices with zero weight, thus extending the results of [6] to the graded case.

1. INTRODUCTION

A complete classification of the nonskewsymmetric solutions of the classical Yang-Baxter equation exists in the case when the underlying structure is a simple Lie algebra; see [1] and [2] for the original proofs by Belavin and Drinfeld, and [5] for a more pedagogical exposition. A similar construction, with natural modifications, works in the super case as well; see [8]. However, it turns out that this may not be easily modified into a full classification result; see [9] for an explicit construction and detailed study of a counterexample.

It is well-known that solutions of the classical Yang-Baxter equation on a Lie algebra give us the semiclassical limits of quantizations on the associated Lie group. In [4], Etingof, Schedler and Schiffmann have explicitly constructed quantizations associated to all solutions coming from the Belavin-Drinfeld result. Their method in fact works for all dynamical r -matrices, i.e. the solutions of the more general dynamical Yang-Baxter equation.

The purpose of this paper is to begin a study of the super analog of the theory of dynamical r -matrices. Ultimately our goal is a full theory of quantum groups in the super setting. We expect that understanding the super solutions of the dynamical Yang-Baxter equation will provide us with valuable insight, and hence help us extend or modify the quantization result cited above to obtain a graded analogue.¹ A possible classification result in this more general setting of dynamical r -matrices may also clarify the so far exceptional case of the classical r -matrices mentioned before.

¹As pointed out by P. Etingof, a different method of quantization, by Etingof and Kazhdan, has been generalized to the super setting, see [7]. However, this is a less constructive technique, and does not fully answer our questions.

2. DYNAMICAL r -MATRICES IN THE SUPER SETTING

2.1. Definitions. Let $\mathfrak{g}$ be a simple Lie superalgebra with non-degenerate Killing form $(\cdot, \cdot)$. Let $\mathfrak{h} \subset \mathfrak{g}$ be a Cartan subsuperalgebra, and let $\Delta \subset \mathfrak{h}^*$ be the set of roots associated to $\mathfrak{h}$. Fix a set of simple roots Γ or equivalently a Borel $\mathfrak{b}$. The *classical dynamical Yang-Baxter equation* for a meromorphic function $r : \mathfrak{h}^* \rightarrow \mathfrak{g} \otimes \mathfrak{g}$ will be:

$$Alt_s(dr) + [r^{12}, r^{13}] + [r^{12}, r^{23}] + [r^{13}, r^{23}] = 0 \quad (1)$$

The differential of r will be defined as above as:

$$\begin{aligned} dr & : \mathfrak{h}^* \longrightarrow \mathfrak{g} \otimes \mathfrak{g} \otimes \mathfrak{g} \\ \lambda & \longmapsto \sum_i x_i \otimes \frac{\partial r}{\partial x_i}(\lambda). \end{aligned}$$

Here $\{x_i\}$ is a basis for $\mathfrak{h}$ so all x_i are even. Recall that $Alt_s : \mathfrak{g}^{\otimes 3} \rightarrow \mathfrak{g}^{\otimes 3}$ is given on homogeneous elements by:

$$Alt_s(a \otimes b \otimes c) = a \otimes b \otimes c + (-1)^{|a|(|b|+|c|)} b \otimes c \otimes a + (-1)^{|c|(|a|+|b|)} c \otimes a \otimes b,$$

In view of all this, we can see that for $r = \sum_i R_{i(1)} \otimes R_{i(2)}$:

$$\begin{aligned} Alt_s(dr) &= \sum_i x_i^{(1)} \left(\frac{\partial r}{\partial x_i} \right)^{(23)} + \sum_i x_i^{(2)} \left(\frac{\partial r}{\partial x_i} \right)^{(31)} \\ &+ \sum_i (-1)^{|R_{i(1)}||R_{i(2)}|} x_i^{(3)} \left(\frac{\partial r}{\partial x_i} \right)^{(12)}. \end{aligned}$$

We will say that a meromorphic function $r : \mathfrak{h}^* \rightarrow \mathfrak{g} \otimes \mathfrak{g}$ is a *super dynamical r -matrix with coupling constant ϵ* if it is a solution to Equation 1 and satisfies the generalized unitarity condition:

$$r(\lambda) + T_s(r)(\lambda) = \epsilon \Omega, \quad (2)$$

where Ω is the Casimir element, i.e. the element of $\mathfrak{g} \otimes \mathfrak{g}$ corresponding to the Killing form. Here, $T_s : V \otimes V \rightarrow V \otimes V$ is the *super twist map* defined on the homogeneous elements of a given super vector space V as

$$T_s(a \otimes b) = (-1)^{|a||b|} b \otimes a.$$

Remark. The above definitions are the natural super analogues of the non-graded terminology used in the study of the dynamical Yang-Baxter equation, see [3] for a survey on the non-graded theory.

2.2. Super Dynamical r -matrices with Zero Weight. A super dynamical r -matrix r satisfies the *zero weight condition* if:

$$[h \otimes 1 + 1 \otimes h, r(\lambda)] = 0 \text{ for all } h \in \mathfrak{h}, \lambda \in \mathfrak{h}^*.$$

In the next two sections we will prove the following two statements:

Theorem 2.1. (1) Let X be a subset of the set of roots Δ of a simple Lie superalgebra $\mathfrak{g}$ with non-degenerate Killing form $(\cdot, \cdot)$ such that:

- (a) If $\alpha, \beta \in X$ and $\alpha + \beta$ is a root, then $\alpha + \beta \in X$, and
- (b) If $\alpha \in X$, then $-\alpha \in X$.

Let $\nu \in \mathfrak{h}^*$, and let $D = \sum_{i < j} D_{ij} dx_i \wedge dx_j$ be a closed meromorphic 2-form on $\mathfrak{h}^*$. If we set $D_{ij} = -D_{ji}$ for $i \geq j$, then the meromorphic function:

$$r(\lambda) = \sum_{i,j=1}^N D_{ij}(\lambda) x_i \otimes x_j + \sum_{\alpha \in X} \frac{(-1)^{|\alpha|} (e_\alpha, e_{-\alpha})}{(\alpha, \lambda - \nu)} e_\alpha \otimes e_{-\alpha}$$

is a super dynamical r -matrix with zero weight and zero coupling constant.

(2) Any super dynamical r -matrix with zero weight and zero coupling constant is of this form.

Theorem 2.2. (1) Let $\mathfrak{g}$ be a simple Lie superalgebra with non-degenerate Killing form $(\cdot, \cdot)$. Let X be a subset of the set of roots Δ of $\mathfrak{g}$ satisfying conditions (a) and (b) of Theorem 2.1. Pick $\nu \in \mathfrak{h}^*$, and define:

$$\varphi_\alpha = \begin{cases} (\epsilon/2) \coth(((-1)^{|\alpha|} (e_\alpha, e_{-\alpha}) (\epsilon/2) (\alpha, \lambda - \nu))) & \text{if } \alpha \in X \\ (\pm \epsilon/2) & \text{if } \alpha \notin X, \text{ negative} \\ \mp (-1)^{|\alpha|} (\epsilon/2) & \text{if } \alpha \notin X, \text{ positive} \end{cases}$$

Let $D = \sum_{i < j} D_{ij} dx_i \wedge dx_j$ be a closed meromorphic 2-form on $\mathfrak{h}^*$. If we set $D_{ij} = -D_{ji}$ for $i \geq j$, then the meromorphic function:

$$r(\lambda) = \sum_{i,j=1}^N D_{ij}(\lambda) x_i \otimes x_j + \frac{\epsilon}{2} \Omega + \sum_{\alpha \in \Delta} \varphi_\alpha e_\alpha \otimes e_{-\alpha}$$

is a super dynamical r -matrix with zero weight and nonzero coupling constant ϵ .

(2) Any super dynamical r -matrix with zero weight and nonzero coupling constant ϵ is of this form.

Remark. Note that if we take the limit as $\epsilon \rightarrow 0$, the above expression reduces to the expression of Theorem 2.1.

2.3. Proof of Theorem 2.1. For any positive root α fix $e_\alpha \in \mathfrak{g}_\alpha$ and pick $e_{-\alpha} \in \mathfrak{g}_{-\alpha}$ dual to e_α i.e.

$$(e_\alpha, e_{-\alpha}) = 1 \text{ for all } \alpha \in \Delta^+.$$

We introduce the following notation:

$$A_\alpha = \begin{cases} (-1)^{|\alpha|} & \text{if } \alpha \text{ is positive} \\ 1 & \text{if } \alpha \text{ is negative} \end{cases}$$

Note that $A_{-\alpha} = (-1)^{|\alpha|} A_{\alpha}$. We can use A_{α} for instance to write the duals of our basis vectors in terms of one another:

$$e_{\alpha}^* = A_{-\alpha} e_{-\alpha}$$

or equivalently:

$$(e_{\alpha}, e_{-\alpha}) = A_{-\alpha}.$$

Let $r : \mathfrak{h}^* \rightarrow \mathfrak{g} \otimes \mathfrak{g}$ be a super dynamical r-matrix with zero coupling constant. Then the zero weight condition on r implies that r has to be of the form:

$$r(\lambda) = \sum_{i,j} D_{ij}(\lambda) x_i \otimes x_j + \sum_{\alpha \in \Delta} \varphi_{\alpha} e_{\alpha} \otimes e_{-\alpha},$$

where D_{ij}, φ_{α} are suitable scalar meromorphic functions such that:

$$D_{ij}(\lambda) = -D_{ji}(\lambda) \quad \text{and} \quad \varphi_{-\alpha} = -(-1)^{|\alpha|} \varphi_{\alpha}$$

One can easily check that there can be no terms mixing the Cartan part with the non-Cartan part, and the conditions on the D_{ij} and the φ_{α} follow from the zero coupling constant.

Equation 1 is an equation in $\mathfrak{g} \otimes \mathfrak{g} \otimes \mathfrak{g}$. The zero coupling constant implies that the left hand side of the equation is skew-symmetric with respect to signed permutations of factors:

$$(12)_s(a \otimes b \otimes c) = (-1)^{|a||b|} b \otimes a \otimes c$$

$$(13)_s(a \otimes b \otimes c) = (-1)^{|a||b|+|a||c|+|b||c|} c \otimes b \otimes a$$

$$(23)_s(a \otimes b \otimes c) = (-1)^{|b||c|} a \otimes c \otimes b$$

Therefore in order to solve Equation 1, it is enough to look at its $\mathfrak{h} \otimes \mathfrak{h} \otimes \mathfrak{h}$, $\mathfrak{h} \otimes \mathfrak{g}_{\alpha} \otimes \mathfrak{g}_{-\alpha}$, and $\mathfrak{g}_{\alpha} \otimes \mathfrak{g}_{\beta} \otimes \mathfrak{g}_{-\alpha-\beta}$ parts.

The $\mathfrak{h} \otimes \mathfrak{h} \otimes \mathfrak{h}$ part is:

$$\begin{aligned} Alt_s(dr) |_{\mathfrak{h} \otimes \mathfrak{h} \otimes \mathfrak{h}} &= \sum_{i,j,k} x_i^{(1)} \frac{\partial D_{jk} x_j x_k^{(23)}}{\partial x_i} + \sum_{i,j,k} x_i^{(2)} \frac{\partial D_{jk} x_j x_k^{(31)}}{\partial x_i} \\ &\quad + \sum_{i,j,k} x_i^{(3)} \frac{\partial D_{jk} x_j x_k^{(12)}}{\partial x_i} \\ &= \sum_{i,j,k} \frac{\partial D_{jk}}{\partial x_i} x_i \otimes x_j \otimes x_k + \sum_{i,j,k} \frac{\partial D_{jk}}{\partial x_i} x_k \otimes x_i \otimes x_j \\ &\quad + \sum_{i,j,k} \frac{\partial D_{jk}}{\partial x_i} x_j \otimes x_k \otimes x_i \\ &= \sum_{i,j,k} \left(\frac{\partial D_{ij}}{\partial x_k} + \frac{\partial D_{jk}}{\partial x_i} + \frac{\partial D_{ki}}{\partial x_j} \right) x_i \otimes x_j \otimes x_k, \end{aligned}$$

and it vanishes if and only if $D = \sum_{i < j} D_{ij} dx_i \wedge dx_j$ is a closed 2-form.

To find the $\mathfrak{h} \otimes \mathfrak{g}_\alpha \otimes \mathfrak{g}_{-\alpha}$ part, we first look at $[[r, r]]$:

$$\begin{aligned} [r^{12}, r^{13}] &= \sum_{i,j,\beta} D_{ij} \varphi_\beta [x_i, e_\beta] \otimes x_j \otimes e_{-\beta} + \sum_{k,l,\alpha} D_{kl} \varphi_\alpha [e_\alpha, x_k] \otimes e_{-\alpha} \otimes x_l \\ &\quad + \sum_{\alpha,\beta} (-1)^{|\alpha||\beta|} \varphi_\alpha \varphi_\beta [e_\alpha, e_\beta] \otimes e_{-\alpha} \otimes e_{-\beta} \end{aligned}$$

$$\begin{aligned} [r^{12}, r^{23}] &= \sum_{i,j,\beta} D_{ij} \varphi_\beta x_i \otimes [x_j, e_\beta] \otimes e_{-\beta} + \sum_{k,l,\alpha} D_{kl} \varphi_\alpha e_\alpha \otimes [e_{-\alpha}, x_k] \otimes x_l \\ &\quad + \sum_{\alpha,\beta} \varphi_\alpha \varphi_\beta e_\alpha \otimes [e_{-\alpha}, e_\beta] \otimes e_{-\beta} \end{aligned}$$

$$\begin{aligned} [r^{13}, r^{23}] &= \sum_{i,j,\beta} D_{ij} \varphi_\beta x_i \otimes e_\beta \otimes [x_j, e_{-\beta}] + \sum_{k,l,\alpha} D_{kl} \varphi_\alpha e_\alpha \otimes x_k \otimes [e_{-\alpha}, x_l] \\ &\quad + \sum_{\alpha,\beta} (-1)^{|\alpha||\beta|} \varphi_\alpha \varphi_\beta e_\alpha \otimes e_\beta \otimes [e_{-\alpha}, e_{-\beta}] \end{aligned}$$

We can now easily see that the $\mathfrak{h} \otimes \mathfrak{g}_\alpha \otimes \mathfrak{g}_{-\alpha}$ part will be:

$$\begin{aligned} Alt_s(dr) \big|_{\mathfrak{h} \otimes \mathfrak{g}_\alpha \otimes \mathfrak{g}_{-\alpha}} &+ \sum_{\alpha,\beta} (-1)^{|\alpha||\beta|} \delta_{\alpha,-\beta} \varphi_\alpha \varphi_\beta [e_\alpha, e_\beta] \otimes e_{-\alpha} \otimes e_{-\beta} + \\ &\sum_{i,j,\beta} D_{ij} \varphi_\beta x_i \otimes [x_j, e_\beta] \otimes e_{-\beta} + \sum_{i,j,\beta} D_{ij} \varphi_\beta x_i \otimes e_\beta \otimes [x_j, e_{-\beta}] \end{aligned}$$

We note that the last two sums cancel out, because:

$$e_\beta \otimes [x_j, e_{-\beta}] = (-\beta)(x_j) e_\beta \otimes e_{-\beta} = -(\beta(x_j)) e_\beta \otimes e_{-\beta} = -[x_j, e_\beta] \otimes e_\beta$$

and we have:

$$\begin{aligned} &Alt_s(dr) \big|_{\mathfrak{h} \otimes \mathfrak{g}_\alpha \otimes \mathfrak{g}_{-\alpha}} + \sum_{\alpha,\beta} (-1)^{|\alpha||\beta|} \delta_{\alpha,-\beta} \varphi_\alpha \varphi_\beta [e_\alpha, e_\beta] \otimes e_{-\alpha} \otimes e_{-\beta} \\ &= \sum_{i,\alpha} \frac{\partial \varphi_\alpha}{\partial x_i} x_i \otimes e_\alpha \otimes e_{-\alpha} + \sum_{\alpha} (-1)^{|\alpha|} \varphi_\alpha \varphi_{-\alpha} [e_\alpha, e_{-\alpha}] \otimes e_{-\alpha} \otimes e_\alpha \\ &= \sum_{i,\alpha} \frac{\partial \varphi_\alpha}{\partial x_i} x_i \otimes e_\alpha \otimes e_{-\alpha} + \sum_{\alpha} (-1)^{|\alpha|} \varphi_\alpha \varphi_{-\alpha} [e_{-\alpha}, e_\alpha] \otimes e_\alpha \otimes e_{-\alpha} \\ &= \sum_{i,\alpha} \frac{\partial \varphi_\alpha}{\partial x_i} x_i \otimes e_\alpha \otimes e_{-\alpha} + \sum_{\alpha} (-1)^{|\alpha|} \varphi_\alpha \varphi_{-\alpha} (-A_\alpha) h_\alpha \otimes e_\alpha \otimes e_{-\alpha} \\ &= \sum_{i,\alpha} \frac{\partial \varphi_\alpha}{\partial x_i} x_i \otimes e_\alpha \otimes e_{-\alpha} + \sum_{\alpha} A_\alpha \varphi_\alpha \varphi_\alpha h_\alpha \otimes e_\alpha \otimes e_{-\alpha} \end{aligned}$$

which we can rewrite as:

$$\sum_{\alpha \in \Delta} \left(\sum_i \frac{\partial \varphi_\alpha}{\partial x_i} x_i + A_\alpha \varphi_\alpha^2 h_\alpha \right) \otimes e_\alpha \otimes e_{-\alpha}$$

where $h_\alpha \in \mathfrak{h}$ is defined by:

$$[e_\alpha, e_{-\alpha}] = (e_\alpha, e_{-\alpha}) h_\alpha = A_{-\alpha} h_\alpha.$$

(Recall that both $\{x_i\}$ and $\{h_\alpha | \alpha \in \Gamma\}$ are bases for $\mathfrak{h}$ and equivalently linear coordinate systems for $\mathfrak{h}^*$). For this term to vanish we must have, for all $\alpha \in \Delta$:

$$\sum_i \frac{\partial \varphi_\alpha}{\partial x_i} x_i + A_\alpha \varphi_\alpha^2 h_\alpha = 0.$$

We can rewrite this as:

$$d\varphi_\alpha + A_\alpha \varphi_\alpha^2 dh_\alpha = 0.$$

If we define μ_α by:

$$\mu_\alpha = \begin{cases} \sqrt{-1} & \text{if } \alpha \text{ is an odd positive root} \\ 1 & \text{otherwise} \end{cases}$$

then $\mu_\alpha^2 = A_\alpha$, and the equation we need to solve is:

$$d\varphi_\alpha + \mu_\alpha^2 \varphi_\alpha^2 dh_\alpha = 0.$$

We assume $\varphi_\alpha \neq 0$ and let $u_\alpha = \mu_\alpha \varphi_\alpha$. Separating variables to integrate we obtain:

$$\frac{1}{\mu_\alpha} \int \frac{-du_\alpha}{u_\alpha^2} = \int dh_\alpha \Rightarrow u_\alpha = \mu_\alpha \varphi_\alpha = \frac{1}{\mu_\alpha h_\alpha + C}$$

and we get:

$$\varphi_\alpha = \frac{A_\alpha}{h_\alpha - \nu_\alpha}$$

for some $\nu_\alpha \in \mathbb{C}$. Here h_α is viewed as a linear function on $\mathfrak{h}^*$ via $h_\alpha(\lambda) = (\alpha, \lambda)$.

Finally we look at the $\mathfrak{g}_\alpha \otimes \mathfrak{g}_\beta \otimes \mathfrak{g}_{-\alpha-\beta}$ part of Equation 1. There is no contribution from the dynamical part; the only terms we need to look at are:

$$\begin{aligned} \sum_{\alpha, \beta} (-1)^{|\alpha||\beta|} \varphi_\alpha \varphi_\beta [e_\alpha, e_\beta] \otimes e_{-\alpha} \otimes e_{-\beta} + \sum_{\alpha, \beta} \varphi_\alpha \varphi_\beta e_\alpha \otimes [e_{-\alpha}, e_\beta] \otimes e_{-\beta} \\ + \sum_{\alpha, \beta} (-1)^{|\alpha||\beta|} \varphi_\alpha \varphi_\beta e_\alpha \otimes e_\beta \otimes [e_{-\alpha}, e_{-\beta}] \end{aligned}$$

Denote by C the matrix of coefficients determined by:

$$[e_\alpha, e_\beta] = C_{\alpha, \beta}^\gamma e_\gamma.$$

Then we can rewrite the terms we are interested in as:

$$\sum_{\alpha, \beta, \gamma} (-1)^{|\alpha||\beta|} \varphi_\alpha \varphi_\beta C_{\alpha, \beta}^\gamma e_\gamma \otimes e_{-\alpha} \otimes e_{-\beta}$$

$$\begin{aligned}
& + \sum_{\alpha,\beta,\gamma} \varphi_\alpha \varphi_\beta C_{-\alpha,\beta}^\gamma e_\alpha \otimes e_\gamma \otimes e_{-\beta} \\
& + \sum_{\alpha,\beta,\gamma} (-1)^{|\alpha||\beta|} \varphi_\alpha \varphi_\beta C_{-\alpha,-\beta}^\gamma e_\alpha \otimes e_\beta \otimes e_\gamma \\
& = \sum_{\alpha,\beta} (-1)^{|\alpha||\beta|} \varphi_\alpha \varphi_\beta C_{\alpha,\beta}^{\alpha+\beta} e_{\alpha+\beta} \otimes e_{-\alpha} \otimes e_{-\beta} \\
& \quad + \sum_{\alpha,\beta} \varphi_\alpha \varphi_\beta C_{-\alpha,\beta}^{-\alpha+\beta} e_\alpha \otimes e_{-\alpha+\beta} \otimes e_{-\beta} \\
& \quad + \sum_{\alpha,\beta} (-1)^{|\alpha||\beta|} \varphi_\alpha \varphi_\beta C_{-\alpha,-\beta}^{-\alpha-\beta} e_\alpha \otimes e_\beta \otimes e_{-\alpha-\beta} \\
& = \sum_{\alpha,\beta} (-1)^{|\alpha||\beta|} \varphi_\alpha \varphi_\beta C_{\alpha,\beta}^{\alpha+\beta} e_{\alpha+\beta} \otimes e_{-\alpha} \otimes e_{-\beta} \\
& \quad + \sum_{\alpha,\beta} \varphi_\alpha \varphi_\beta C_{-\alpha,\beta}^{-\alpha+\beta} e_\alpha \otimes e_{-\alpha+\beta} \otimes e_{-\beta} \\
& \quad + \sum_{\alpha,\beta} (-1)^{|\alpha||\beta|} \varphi_\alpha \varphi_\beta C_{-\alpha,-\beta}^{-\alpha-\beta} e_\alpha \otimes e_\beta \otimes e_{-\alpha-\beta}
\end{aligned}$$

We want the coefficient in front of the term $e_\alpha \otimes e_\beta \otimes e_{-\alpha-\beta}$ to vanish:

$$\begin{aligned}
& (-1)^{|\beta||\alpha+\beta|} \varphi_{-\beta} \varphi_{\alpha+\beta} C_{-\beta,\alpha+\beta}^\alpha + \varphi_\alpha \varphi_{\alpha+\beta} C_{-\alpha,\alpha+\beta}^\beta \\
& \quad + (-1)^{|\alpha||\beta|} \varphi_\alpha \varphi_\beta C_{-\alpha,-\beta}^{-\alpha-\beta} = \\
& -(-1)^{|\alpha||\beta|} \varphi_\beta \varphi_{\alpha+\beta} C_{-\beta,\alpha+\beta}^\alpha + \varphi_\alpha \varphi_{\alpha+\beta} C_{-\alpha,\alpha+\beta}^\beta \\
& \quad + (-1)^{|\alpha||\beta|} \varphi_\alpha \varphi_\beta C_{-\alpha,-\beta}^{-\alpha-\beta} = 0
\end{aligned}$$

We now compute these constants:

$$\begin{aligned}
C_{-\beta,\alpha+\beta}^\alpha &= ([e_{-\beta}, e_{\alpha+\beta}], e_\alpha^*) = ([e_{-\beta}, e_{\alpha+\beta}], A_{-\alpha} e_{-\alpha}) \\
&= \frac{A_{-\alpha}}{C_{\alpha,\beta}^{\alpha+\beta}} ([e_{-\beta}, [e_\alpha, e_\beta]], e_{-\alpha}) \\
&= \frac{-(-1)^{|\alpha||\beta|} A_{-\alpha}}{C_{\alpha,\beta}^{\alpha+\beta}} ([e_{-\beta}, [e_\beta, e_\alpha]], e_{-\alpha}) \\
&= \frac{-(-1)^{|\alpha||\beta|} A_{-\alpha}}{C_{\alpha,\beta}^{\alpha+\beta}} ([e_{-\beta}, e_\beta], e_\alpha + (-1)^{|\beta|} [e_\beta, [e_{-\beta}, e_\alpha]], e_{-\alpha}) \\
&= \frac{-(-1)^{|\alpha||\beta|} A_{-\alpha}}{C_{\alpha,\beta}^{\alpha+\beta}} ([e_{-\beta}, e_\beta], e_\alpha, e_{-\alpha}) \\
&= \frac{(-1)^{|\alpha||\beta|} (-1)^{|\beta|} A_{-\alpha}}{C_{\alpha,\beta}^{\alpha+\beta}} ([e_\beta, e_{-\beta}], e_\alpha, e_{-\alpha}) \\
&= \frac{(-1)^{|\alpha||\beta|} (-1)^{|\beta|} A_{-\alpha} A_{-\beta}}{C_{\alpha,\beta}^{\alpha+\beta}} (h_\beta, [e_\alpha, e_{-\alpha}])
\end{aligned}$$

$$= \frac{(-1)^{|\alpha||\beta|}(-1)^{|\beta|}A_{-\beta}}{C_{\alpha,\beta}^{\alpha+\beta}}(h_\alpha, h_\beta)$$

where we use the Jacobi identity and $(A_\alpha)^2 = 1$. Similar computations yield:

$$\begin{aligned} C_{-\alpha,\alpha+\beta}^\beta &= \frac{-(-1)^{|\alpha|}A_{-\alpha}}{C_{\alpha,\beta}^{\alpha+\beta}}(h_\alpha, h_\beta) \\ C_{-\alpha,-\beta}^{-\alpha-\beta} &= \frac{(-1)^{|\alpha||\beta|}(-1)^{|\alpha|+|\beta|}A_{\alpha+\beta}A_{-\alpha}A_{-\beta}}{C_{\alpha,\beta}^{\alpha+\beta}}(h_\alpha, h_\beta) \end{aligned}$$

Recall that if $\alpha + \beta$ is a root, $(h_\alpha, h_\beta) \neq 0$ and $C_{\alpha,\beta}^{\alpha+\beta} \neq 0$. Then we can rewrite the coefficient of $e_\alpha \otimes e_\beta \otimes e_{-\alpha-\beta}$:

$$\begin{aligned} 0 &= -(-1)^{|\alpha||\beta|}\varphi_\beta\varphi_{\alpha+\beta}C_{-\beta,\alpha+\beta}^\alpha + \varphi_\alpha\varphi_{\alpha+\beta}C_{-\alpha,\alpha+\beta}^\beta \\ &\quad + (-1)^{|\alpha||\beta|}\varphi_\alpha\varphi_\beta C_{-\alpha,-\beta}^{-\alpha-\beta} \\ 0 &= -(-1)^{|\alpha||\beta|}\varphi_\beta\varphi_{\alpha+\beta}\frac{(-1)^{|\alpha||\beta|}(-1)^{|\beta|}A_{-\beta}}{C_{\alpha,\beta}^{\alpha+\beta}}(h_\alpha, h_\beta) \\ &\quad + \varphi_\alpha\varphi_{\alpha+\beta}\frac{-(-1)^{|\alpha|}A_{-\alpha}}{C_{\alpha,\beta}^{\alpha+\beta}}(h_\alpha, h_\beta) \\ &\quad + (-1)^{|\alpha||\beta|}\varphi_\alpha\varphi_\beta\frac{(-1)^{|\alpha||\beta|}(-1)^{|\alpha|+|\beta|}A_{\alpha+\beta}A_{-\alpha}A_{-\beta}}{C_{\alpha,\beta}^{\alpha+\beta}}(h_\alpha, h_\beta) \\ 0 &= -(-1)^{|\alpha||\beta|}\varphi_\beta\varphi_{\alpha+\beta}(-1)^{|\alpha||\beta|}(-1)^{|\beta|}A_{-\alpha} + \varphi_\alpha\varphi_{\alpha+\beta}(-(-1)^{|\alpha|})A_{-\beta} \\ &\quad + (-1)^{|\alpha||\beta|}\varphi_\alpha\varphi_\beta(-1)^{|\alpha||\beta|}(-1)^{|\alpha|+|\beta|}A_{\alpha+\beta} \\ 0 &= -(-1)^{|\beta|}\varphi_\beta\varphi_{\alpha+\beta}A_{-\alpha} - (-1)^{|\alpha|}\varphi_\alpha\varphi_{\alpha+\beta}A_{-\beta} \\ &\quad + (-1)^{|\alpha|+|\beta|}\varphi_\alpha\varphi_\beta A_{\alpha+\beta} \end{aligned}$$

which can be rewritten as:

$$\begin{aligned} A_{\alpha+\beta}\varphi_\alpha\varphi_\beta &= A_\alpha\varphi_\beta\varphi_{\alpha+\beta} + A_\beta\varphi_\alpha\varphi_{\alpha+\beta} \\ &= (A_\alpha\varphi_\beta + A_\beta\varphi_\alpha)\varphi_{\alpha+\beta} \end{aligned}$$

where we use $A_\alpha = (-1)^{|\alpha|}A_{-\alpha}$ and $(A_\alpha)^2 = 1$.

Define the set

$$X = \{\alpha \in \Delta \mid \varphi_\alpha \neq 0\}$$

Then it is easy to see that X is closed under addition and changing signs. In other words, if $\alpha, \beta \in X$, then so are $\alpha + \beta$ and $-\alpha$. These follow directly from the above equation relating φ_α and φ_β to $\varphi_{\alpha+\beta}$, and the unitarity property (i.e. $r = -T_s(r)$).

Next assume α and β are positive and in X . Then the above calculations yield:

$$\begin{aligned} A_{\alpha+\beta}\varphi_\alpha\varphi_\beta &= (A_\alpha\varphi_\beta + A_\beta\varphi_\alpha)\varphi_{\alpha+\beta} \\ (-1)^{|\alpha|+|\beta|}\varphi_\alpha\varphi_\beta &= ((-1)^{|\alpha|}\varphi_\beta + (-1)^{|\beta|}\varphi_\alpha)\varphi_{\alpha+\beta} \\ (-1)^{|\alpha|+|\beta|}\left(\frac{(-1)^{|\alpha|}}{h_\alpha - \nu_\alpha}\right)\left(\frac{(-1)^{|\beta|}}{h_\beta - \nu_\beta}\right) &= \left(\frac{(-1)^{|\alpha+\beta|}}{h_\beta - \nu_\beta} + \frac{(-1)^{|\alpha+\beta|}}{h_\alpha - \nu_\alpha}\right)\frac{(-1)^{|\alpha+\beta|}}{h_{\alpha+\beta} - \nu_{\alpha+\beta}} \\ \left(\frac{1}{h_\alpha - \nu_\alpha}\right)\left(\frac{1}{h_\beta - \nu_\beta}\right) &= \left(\frac{1}{h_\beta - \nu_\beta} + \frac{1}{h_\alpha - \nu_\alpha}\right)\frac{1}{h_{\alpha+\beta} - \nu_{\alpha+\beta}} \end{aligned}$$

Since $h_{\alpha+\beta} = h_\alpha + h_\beta$ we must have:

$$\nu_{\alpha+\beta} = \nu_\alpha + \nu_\beta$$

Also recall that $\varphi_{-\alpha} = -(-1)^{|\alpha|}\varphi_\alpha$ and so:

$$\frac{A_{-\alpha}}{h_{-\alpha} - \nu_{-\alpha}} = -(-1)^{|\alpha|}\frac{A_\alpha}{h_\alpha - \nu_\alpha}$$

which implies that $\nu_{-\alpha} = -\nu_\alpha$. Therefore we can conclude that there is some $\nu \in \mathfrak{h}^*$ such that $\nu_\alpha = (\alpha, \nu)$ for all $\alpha \in X$. This completes the proof of the theorem. ■

2.4. Proof of Theorem 2.2. We start with fixing a basis for the non-Cartan part of $\mathfrak{g}$ in the same manner as above. In other words for any positive root α we fix $e_\alpha \in \mathfrak{g}_\alpha$ and pick $e_{-\alpha} \in \mathfrak{g}_{-\alpha}$ dual to e_α i.e.

$$(e_\alpha, e_{-\alpha}) = 1 \text{ for all } \alpha \in \Delta^+.$$

We again need the following notation:

$$A_\alpha = \begin{cases} (-1)^{|\alpha|} & \text{if } \alpha \text{ is positive} \\ 1 & \text{if } \alpha \text{ is negative} \end{cases}$$

and we note once again that $A_{-\alpha} = (-1)^{|\alpha|}A_\alpha$.

Let $r : \mathfrak{h}^* \rightarrow \mathfrak{g} \otimes \mathfrak{g}$ be a meromorphic map, $\Omega \in \mathfrak{g} \otimes \mathfrak{g}$ the Casimir element, and ϵ a nonzero complex number. Introduce a second meromorphic function $s : \mathfrak{h}^* \rightarrow \mathfrak{g} \otimes \mathfrak{g}$ by:

$$s(\lambda) = r(\lambda) - \frac{\epsilon}{2}\Omega \quad \text{for all } \lambda \in \mathfrak{h}^*.$$

We will now prove the following technical lemma:

Lemma 2.1. *The map r is a super dynamical r -matrix with zero weight and coupling constant ϵ if and only if s satisfies the zero weight condition:*

$$[h \otimes 1 + 1 \otimes h, s(\lambda)] = 0 \text{ for all } h \in \mathfrak{h}, \lambda \in \mathfrak{h}^*,$$

the unitarity condition:

$$s(\lambda) + T_s(s)(\lambda) = 0,$$

and the following modified version of the dynamical Yang-Baxter equation:

$$Alt_s(ds) + [[s, s]] + \frac{\epsilon^2}{4} [[\Omega, \Omega]] = 0, \quad (3)$$

where, for any 2-tensor r :

$$[[r, r]] = [r^{12}, r^{13}] + [r^{12}, r^{23}] + [r^{13}, r^{23}].$$

Remark. This is exactly Lemma 3.9 of [6]. Its proof consists mainly of a direct calculation, but the computations are significantly more involved in the super case. In any case, the proof will be given here for completeness.

Proof: The zero weight condition and the unitarity condition on s imply that s has to be of the form:

$$s(\lambda) = \sum_{i,j} D_{ij}(\lambda) h_i \otimes h_j + \sum_{\alpha \in \Delta} \varphi_\alpha e_\alpha \otimes e_{-\alpha},$$

where $\{h_i\}$ is a basis for $\mathfrak{h}$, and D_{ij}, φ_α are suitable scalar meromorphic functions such that:

$$D_{ij}(\lambda) = -D_{ji}(\lambda) \quad \text{and} \quad \varphi_{-\alpha} = -(-1)^{|\alpha|} \varphi_\alpha$$

One can easily check that there can be no terms mixing the Cartan part with the non-Cartan part, and the conditions on the D_{ij} and the φ_α follow from unitarity.

We have:

$$[[r, r]] = \left[\left[s + \frac{\epsilon}{2} \Omega, s + \frac{\epsilon}{2} \Omega \right] \right]$$

so it suffices to prove that $[[r, r]] - [[s, s]] - \frac{\epsilon^2}{4} [[\Omega, \Omega]]$ is zero. In other words we need to show that

$$\frac{\epsilon}{2} ([s^{12}, \Omega^{13}] + [\Omega^{12}, s^{13}] + [s^{12}, \Omega^{23}] + [\Omega^{12}, s^{23}] + [s^{13}, \Omega^{23}] + [\Omega^{13}, s^{23}])$$

vanishes.

If $\{h^i\}$ is the dual basis in $\mathfrak{h}$ to $\{h_i\}$, then we can write the Casimir element as:

$$\Omega = \sum_i h_i \otimes h^i + \sum_{\alpha \in \Delta} A_\alpha e_\alpha \otimes e_{-\alpha}$$

and we have:

$$\begin{aligned}
[s^{12}, \Omega^{13}] &= \sum_{i,j,k} D_{ij} [h_i, h_k] \otimes h_j \otimes h^k + \sum_{\alpha,k} \varphi_\alpha [e_\alpha, h_k] \otimes e_{-\alpha} \otimes h^k \\
&+ \sum_{i,j,\beta} D_{ij} A_\beta [h_i, e_\beta] \otimes h_j \otimes e_{-\beta} \\
&\quad + \sum_{\alpha,\beta} (-1)^{|\alpha||\beta|} \varphi_\alpha A_\beta [e_\alpha, e_\beta] \otimes e_{-\alpha} \otimes e_{-\beta} \\
[\Omega^{12}, s^{13}] &= \sum_{i,j,k} D_{jk} [h_i, h_j] \otimes h^i \otimes h_k + \sum_{i,\beta} \varphi_\beta [h_i, e_\beta] \otimes h^i \otimes e_{-\beta} \\
&+ \sum_{\alpha,j,k} A_\alpha D_{jk} [e_\alpha, h_j] \otimes e_{-\alpha} \otimes h_k \\
&\quad + \sum_{\alpha,\beta} (-1)^{|\alpha||\beta|} A_\alpha \varphi_\beta [e_\alpha, e_\beta] \otimes e_{-\alpha} \otimes e_{-\beta} \\
[s^{12}, \Omega^{23}] &= \sum_{i,j,k} D_{ij} h_i \otimes [h_j, h_k] \otimes h^k + \sum_{\alpha,k} \varphi_\alpha e_\alpha \otimes [e_{-\alpha}, h_k] \otimes h^k \\
&+ \sum_{i,j,\beta} D_{ij} A_\beta h_i \otimes [h_j, e_\beta] \otimes e_{-\beta} \\
&\quad + \sum_{\alpha,\beta} \varphi_\alpha A_\beta e_\alpha \otimes [e_{-\alpha}, e_\beta] \otimes e_{-\beta} \\
[\Omega^{12}, s^{23}] &= \sum_{i,j,k} D_{jk} h_i \otimes [h^i, h_j] \otimes h_k + \sum_{i,\beta} \varphi_\beta h_i \otimes [h^i, e_\beta] \otimes e_{-\beta} \\
&+ \sum_{\alpha,j,k} A_\alpha D_{jk} e_\alpha \otimes [e_{-\alpha}, h_j] \otimes h_k \\
&\quad + \sum_{\alpha,\beta} A_\alpha \varphi_\beta e_\alpha \otimes [e_{-\alpha}, e_\beta] \otimes e_{-\beta} \\
[s^{13}, \Omega^{23}] &= \sum_{i,j,k} D_{ij} h_i \otimes h_k \otimes [h_j, h^k] + \sum_{\alpha,k} \varphi_\alpha e_\alpha \otimes h_k \otimes [e_{-\alpha}, h^k] \\
&+ \sum_{i,j,\beta} D_{ij} A_\beta h_i \otimes e_\beta \otimes [h_j, e_{-\beta}] \\
&\quad + \sum_{\alpha,\beta} (-1)^{|\alpha||\beta|} \varphi_\alpha A_\beta e_\alpha \otimes e_\beta \otimes [e_{-\alpha}, e_{-\beta}] \\
[\Omega^{13}, s^{23}] &= \sum_{i,j,k} D_{jk} h_i \otimes h_j \otimes [h^i, h_k] + \sum_{i,\beta} \varphi_\beta h_i \otimes e_\beta \otimes [h^i, e_{-\beta}] \\
&+ \sum_{\alpha,j,k} A_\alpha D_{jk} e_\alpha \otimes h_j \otimes [e_{-\alpha}, h_k] \\
&\quad + \sum_{\alpha,\beta} (-1)^{|\alpha||\beta|} A_\alpha \varphi_\beta e_\alpha \otimes e_\beta \otimes [e_{-\alpha}, e_{-\beta}]
\end{aligned}$$

Clearly the first sum in each of these six terms is zero, as the h_i are all in the Cartan. It is also easy to see that the sum of all the third sums vanishes as well:

$$\begin{aligned}
\sum_{i,j,\beta} D_{ij} A_\beta [h_i, e_\beta] \otimes h_j \otimes e_{-\beta} &= \sum_{i,j,\beta} D_{ij} A_\beta \beta(h_i) e_\beta \otimes h_j \otimes e_{-\beta} \\
&= - \sum_{i,j,\beta} (-D_{ij}) A_\beta e_\beta \otimes h_j \otimes [e_{-\beta}, h_i] \\
&= - \sum_{\alpha,j,k} D_{j,k} A_\alpha e_\alpha \otimes h_j \otimes [e_{-\alpha}, h_k]
\end{aligned}$$

where we use $D_{ij} = -D_{ji}$,

$$\begin{aligned}
\sum_{\alpha,j,k} A_\alpha D_{jk} [e_\alpha, h_j] \otimes e_{-\alpha} \otimes h_k &= - \sum_{\alpha,j,k} A_\alpha D_{jk} \alpha(h_j) e_\alpha \otimes e_{-\alpha} \otimes h_k \\
&= - \sum_{\alpha,j,k} A_\alpha D_{jk} e_\alpha \otimes [e_{-\alpha}, h_j] \otimes h_k,
\end{aligned}$$

and:

$$\begin{aligned}
\sum_{i,j,\beta} D_{ij} A_\beta h_i \otimes [h_j, e_\beta] \otimes e_{-\beta} &= \sum_{i,j,\beta} D_{ij} A_\beta \beta(h_j) h_i \otimes e_\beta \otimes e_{-\beta} \\
&= - \sum_{i,j,\beta} D_{ij} A_\beta h_i \otimes e_\beta \otimes [h_j, e_{-\beta}].
\end{aligned}$$

We can also see that:

$$\begin{aligned}
\sum_{\alpha,k} \varphi_\alpha [e_\alpha, h_k] \otimes e_{-\alpha} \otimes h^k &= \sum_{\alpha,k} \varphi_\alpha (-\alpha(h_k) e_\alpha) \otimes e_{-\alpha} \otimes h^k \\
&= - \sum_{\alpha,k} \varphi_\alpha e_\alpha \otimes [e_{-\alpha}, h_k] \otimes h^k,
\end{aligned}$$

and:

$$\begin{aligned}
\sum_{i,\beta} \varphi_\beta h_i \otimes [h^i, e_\beta] \otimes e_{-\beta} &= \sum_{i,\beta} \varphi_\beta \beta(h^i) h_i \otimes e_\beta \otimes e_{-\beta} \\
&= - \sum_{i,\beta} \varphi_\beta h_i \otimes e_\beta \otimes [h^i, e_{-\beta}]
\end{aligned}$$

but:

$$\begin{aligned}
\sum_{i,\beta} \varphi_\beta [h_i, e_\beta] \otimes h^i \otimes e_{-\beta} &= \sum_{i,\beta} \varphi_\beta \beta(h_i) e_\beta \otimes h^i \otimes e_{-\beta} \\
&= \sum_{\beta} \varphi_\beta e_\beta \otimes \left(\sum_i \beta(h_i) h^i \right) \otimes e_{-\beta} \\
&= \sum_{\beta} \varphi_\beta e_\beta \otimes h_\beta \otimes e_{-\beta}
\end{aligned}$$

$$\begin{aligned}
 \sum_{\alpha,k} \varphi_\alpha e_\alpha \otimes h_k \otimes [e_{-\alpha}, h^k] &= \sum_{\alpha,k} \varphi_\alpha \alpha(h^k) e_\alpha \otimes h_k \otimes e_{-\alpha} \\
 &= \sum_{\alpha} \varphi_\alpha e_\alpha \otimes \left(\sum_k \alpha(h^k) h_k \right) \otimes e_{-\alpha} \\
 &= \sum_{\alpha} \varphi_\alpha e_\alpha \otimes h_\alpha \otimes e_{-\alpha}
 \end{aligned}$$

where $h_\alpha \in \mathfrak{h}$ is defined as usual by $A_{-\alpha} h_\alpha = (e_\alpha, e_{-\alpha}) h_\alpha = [e_\alpha, e_{-\alpha}]$.

Next we look at the six remaining terms, the fourth sum in each term above. We note that:

$$\begin{aligned}
 \sum_{\alpha,\beta} (-1)^{|\alpha||\beta|} \varphi_\alpha A_\beta [e_\alpha, e_\beta] \otimes e_{-\alpha} &\otimes e_{-\beta} \delta_{\alpha,-\beta} \\
 &= \sum_{\alpha} (-1)^{|\alpha|} \varphi_\alpha A_{-\alpha} [e_\alpha, e_{-\alpha}] \otimes e_{-\alpha} \otimes e_\alpha \\
 \sum_{\alpha,\beta} (-1)^{|\alpha||\beta|} A_\alpha \varphi_\beta [e_\alpha, e_\beta] \otimes e_{-\alpha} &\otimes e_{-\beta} \delta_{\alpha,-\beta} \\
 &= \sum_{\alpha} (-1)^{|\alpha|} A_\alpha \varphi_{-\alpha} [e_\alpha, e_{-\alpha}] \otimes e_{-\alpha} \otimes e_\alpha \\
 \sum_{\alpha,\beta} \varphi_\alpha A_\beta e_\alpha \otimes [e_{-\alpha}, e_\beta] &\otimes e_{-\beta} \delta_{\alpha,\beta} \\
 &= \sum_{\alpha} \varphi_\alpha A_\alpha e_\alpha \otimes [e_{-\alpha}, e_\alpha] \otimes e_{-\alpha} \\
 \sum_{\alpha,\beta} A_\alpha \varphi_\beta e_\alpha \otimes [e_{-\alpha}, e_\beta] &\otimes e_{-\beta} \delta_{\alpha,\beta} \\
 &= \sum_{\alpha} A_\alpha \varphi_\alpha e_\alpha \otimes [e_{-\alpha}, e_\alpha] \otimes e_{-\alpha} \\
 \sum_{\alpha,\beta} (-1)^{|\alpha||\beta|} \varphi_\alpha A_\beta e_\alpha \otimes e_\beta &\otimes [e_{-\alpha}, e_{-\beta}] \delta_{\alpha,-\beta} \\
 &= \sum_{\alpha} (-1)^{|\alpha|} \varphi_\alpha A_{-\alpha} e_\alpha \otimes e_{-\alpha} \otimes [e_{-\alpha}, e_\alpha] \\
 \sum_{\alpha,\beta} (-1)^{|\alpha||\beta|} A_\alpha \varphi_\beta e_\alpha \otimes e_\beta &\otimes [e_{-\alpha}, e_{-\beta}] \delta_{\alpha,-\beta} \\
 &= \sum_{\alpha} (-1)^{|\alpha|} A_\alpha \varphi_{-\alpha} e_\alpha \otimes e_{-\alpha} \otimes [e_{-\alpha}, e_\alpha].
 \end{aligned}$$

We can see that the first two and the last two of these cancel out one another because:

$$\varphi_\alpha A_{-\alpha} = (-1)^{|\alpha|} \varphi_\alpha A_\alpha \quad A_\alpha \varphi_{-\alpha} = -(-1)^{|\alpha|} A_\alpha \varphi_\alpha$$

and the center two add up to give:

$$\begin{aligned}
2 \sum_{\alpha} \varphi_{\alpha} A_{\alpha} e_{\alpha} \otimes [e_{-\alpha}, e_{\alpha}] &\otimes e_{-\alpha} \\
&= -2 \sum_{\alpha} \varphi_{\alpha} (-1)^{|\alpha|} A_{\alpha} e_{\alpha} \otimes [e_{\alpha}, e_{-\alpha}] \otimes e_{-\alpha} \\
&= -2 \sum_{\alpha} \varphi_{\alpha} (A_{-\alpha})^2 e_{\alpha} \otimes h_{\alpha} \otimes e_{-\alpha} \\
&= -2 \sum_{\alpha} \varphi_{\alpha} e_{\alpha} \otimes h_{\alpha} \otimes e_{-\alpha}
\end{aligned}$$

because $A_{\alpha}^2 = 1$. This cancels the terms remaining from the second sums:

$$\sum_{\beta} \varphi_{\beta} e_{\beta} \otimes h_{\beta} \otimes e_{-\beta} + \sum_{\alpha} \varphi_{\alpha} e_{\alpha} \otimes h_{\alpha} \otimes e_{-\alpha} = 2 \sum_{\beta} \varphi_{\beta} e_{\beta} \otimes h_{\beta} \otimes e_{-\beta}$$

Finally the only terms remaining that we need to check are the terms of the form $e_{\alpha} \otimes e_{\beta} \otimes e_{\gamma}$ which all come from the fourth sums. If C is again the matrix of coefficients determined by:

$$[e_{\alpha}, e_{\beta}] = \sum_{\gamma} C_{\alpha, \beta}^{\gamma} e_{\gamma} = C_{\alpha, \beta}^{\alpha+\beta} e_{\alpha+\beta},$$

then we can write the last six terms as follows:

$$\begin{aligned}
&\sum_{\alpha, \beta} (-1)^{|\alpha|+|\beta|} \varphi_{\alpha} A_{\beta} [e_{\alpha}, e_{\beta}] \otimes e_{-\alpha} \otimes e_{-\beta} \\
&= \sum_{\alpha, \beta} (-1)^{|\alpha|+|\beta|} \varphi_{\alpha} A_{\beta} C_{\alpha, \beta}^{\alpha+\beta} e_{\alpha+\beta} \otimes e_{-\alpha} \otimes e_{-\beta} \\
&\sum_{\alpha, \beta} (-1)^{|\alpha|+|\beta|} A_{\alpha} \varphi_{\beta} [e_{\alpha}, e_{\beta}] \otimes e_{-\alpha} \otimes e_{-\beta} \\
&= \sum_{\alpha, \beta} (-1)^{|\alpha|+|\beta|} A_{\alpha} \varphi_{\beta} C_{\alpha, \beta}^{\alpha+\beta} e_{\alpha+\beta} \otimes e_{-\alpha} \otimes e_{-\beta} \\
&\sum_{\alpha, \beta} \varphi_{\alpha} A_{\beta} e_{\alpha} \otimes [e_{-\alpha}, e_{\beta}] \otimes e_{-\beta} \\
&= \sum_{\alpha, \beta} \varphi_{\alpha} A_{\beta} C_{-\alpha, \beta}^{-\alpha+\beta} e_{\alpha} \otimes e_{-\alpha+\beta} \otimes e_{-\beta} \\
&\sum_{\alpha, \beta} A_{\alpha} \varphi_{\beta} e_{\alpha} \otimes [e_{-\alpha}, e_{\beta}] \otimes e_{-\beta} \\
&= \sum_{\alpha, \beta} A_{\alpha} \varphi_{\beta} C_{-\alpha, \beta}^{-\alpha+\beta} e_{\alpha} \otimes e_{-\alpha+\beta} \otimes e_{-\beta} \\
&\sum_{\alpha, \beta} (-1)^{|\alpha|+|\beta|} \varphi_{\alpha} A_{\beta} e_{\alpha} \otimes e_{\beta} \otimes [e_{-\alpha}, e_{-\beta}]
\end{aligned}$$

$$\begin{aligned}
 &= \sum_{\alpha, \beta} (-1)^{|\alpha||\beta|} \varphi_\alpha A_\beta C_{-\alpha, -\beta}^{-\alpha-\beta} e_\alpha \otimes e_\beta \otimes e_{-\alpha-\beta} \\
 \sum_{\alpha, \beta} (-1)^{|\alpha||\beta|} A_\alpha \varphi_\beta e_\alpha &\otimes e_\beta \otimes [e_{-\alpha}, e_{-\beta}] \\
 &= \sum_{\alpha, \beta} (-1)^{|\alpha||\beta|} A_\alpha \varphi_\beta C_{-\alpha, -\beta}^{-\alpha-\beta} e_\alpha \otimes e_\beta \otimes e_{-\alpha-\beta}
 \end{aligned}$$

We write down the contribution of each of these six sums to the coefficient in front of $e_\alpha \otimes e_\beta \otimes e_{-\alpha-\beta}$:

$$\begin{aligned}
 \sum_{\alpha, \beta} (-1)^{|\alpha||\beta|} \varphi_\alpha A_\beta C_{\alpha, \beta}^{\alpha+\beta} e_{\alpha+\beta} &\otimes e_{-\alpha} \otimes e_{-\beta} \\
 &: (-1)^{|\beta||\alpha+\beta|} \varphi_{-\beta} A_{\alpha+\beta} C_{-\beta, \alpha+\beta}^\alpha \\
 \sum_{\alpha, \beta} (-1)^{|\alpha||\beta|} A_\alpha \varphi_\beta C_{\alpha, \beta}^{\alpha+\beta} e_{\alpha+\beta} &\otimes e_{-\alpha} \otimes e_{-\beta} \\
 &: (-1)^{|\beta||\alpha+\beta|} A_{-\beta} \varphi_{\alpha+\beta} C_{-\beta, \alpha+\beta}^\alpha \\
 \sum_{\alpha, \beta} \varphi_\alpha A_\beta C_{-\alpha, \beta}^{-\alpha+\beta} e_\alpha &\otimes e_{-\alpha+\beta} \otimes e_{-\beta} \\
 &: \varphi_\alpha A_{\alpha+\beta} C_{-\alpha, \alpha+\beta}^\beta \\
 \sum_{\alpha, \beta} A_\alpha \varphi_\beta C_{-\alpha, \beta}^{-\alpha+\beta} e_\alpha &\otimes e_{-\alpha+\beta} \otimes e_{-\beta} \\
 &: A_\alpha \varphi_{\alpha+\beta} C_{-\alpha, \alpha+\beta}^\beta \\
 \sum_{\alpha, \beta} (-1)^{|\alpha||\beta|} \varphi_\alpha A_\beta C_{-\alpha, -\beta}^{-\alpha-\beta} e_\alpha &\otimes e_\beta \otimes e_{-\alpha-\beta} \\
 &: (-1)^{|\alpha||\beta|} \varphi_\alpha A_\beta C_{-\alpha, -\beta}^{-\alpha-\beta} \\
 \sum_{\alpha, \beta} (-1)^{|\alpha||\beta|} A_\alpha \varphi_\beta C_{-\alpha, -\beta}^{-\alpha-\beta} e_\alpha &\otimes e_\beta \otimes e_{-\alpha-\beta} \\
 &: (-1)^{|\alpha||\beta|} A_\alpha \varphi_\beta C_{-\alpha, -\beta}^{-\alpha-\beta}
 \end{aligned}$$

Gathering like terms together we see that the coefficient we want is:

$$\begin{aligned}
 &(-1)^{|\beta||\alpha+\beta|} \varphi_{-\beta} A_{\alpha+\beta} C_{-\beta, \alpha+\beta}^\alpha + (-1)^{|\beta||\alpha+\beta|} A_{-\beta} \varphi_{\alpha+\beta} C_{-\beta, \alpha+\beta}^\alpha \\
 &+ \varphi_\alpha A_{\alpha+\beta} C_{-\alpha, \alpha+\beta}^\beta + A_\alpha \varphi_{\alpha+\beta} C_{-\alpha, \alpha+\beta}^\beta \\
 &+ (-1)^{|\alpha||\beta|} \varphi_\alpha A_\beta C_{-\alpha, -\beta}^{-\alpha-\beta} + (-1)^{|\alpha||\beta|} A_\alpha \varphi_\beta C_{-\alpha, -\beta}^{-\alpha-\beta}
 \end{aligned}$$

or equivalently:

$$\begin{aligned}
 &= \varphi_\alpha A_{\alpha+\beta} C_{-\alpha, \alpha+\beta}^\beta + (-1)^{|\alpha||\beta|} \varphi_\alpha A_\beta C_{-\alpha, -\beta}^{-\alpha-\beta} \\
 &+ (-1)^{|\beta||\alpha+\beta|} \varphi_{-\beta} A_{\alpha+\beta} C_{-\beta, \alpha+\beta}^\alpha + (-1)^{|\alpha||\beta|} A_\alpha \varphi_\beta C_{-\alpha, -\beta}^{-\alpha-\beta}
 \end{aligned}$$

$$\begin{aligned}
& + A_\alpha \varphi_{\alpha+\beta} C_{-\alpha, \alpha+\beta}^\beta + (-1)^{|\beta||\alpha+\beta|} A_{-\beta} \varphi_{\alpha+\beta} C_{-\beta, \alpha+\beta}^\alpha \\
= & \varphi_\alpha \left(A_{\alpha+\beta} C_{-\alpha, \alpha+\beta}^\beta + (-1)^{|\alpha||\beta|} A_\beta C_{-\alpha, -\beta}^{-\alpha-\beta} \right) \\
& + \varphi_\beta \left(-(-1)^{|\beta|} (-1)^{|\beta||\alpha+\beta|} A_{\alpha+\beta} C_{-\beta, \alpha+\beta}^\alpha + (-1)^{|\alpha||\beta|} A_\alpha C_{-\alpha, -\beta}^{-\alpha-\beta} \right) \\
& + \varphi_{\alpha+\beta} \left(A_\alpha C_{-\alpha, \alpha+\beta}^\beta + (-1)^{|\beta||\alpha+\beta|} A_{-\beta} C_{-\beta, \alpha+\beta}^\alpha \right)
\end{aligned}$$

We have computed earlier that:

$$\begin{aligned}
C_{-\beta, \alpha+\beta}^\alpha &= \frac{(-1)^{|\alpha||\beta|} (-1)^{|\beta|} A_{-\beta}}{C_{\alpha, \beta}^{\alpha+\beta}} (h_\alpha, h_\beta) \\
C_{-\alpha, \alpha+\beta}^\beta &= \frac{-(-1)^{|\alpha|} A_{-\alpha}}{C_{\alpha, \beta}^{\alpha+\beta}} (h_\alpha, h_\beta) \\
C_{-\alpha, -\beta}^{-\alpha-\beta} &= \frac{(-1)^{|\alpha||\beta|} (-1)^{|\alpha|+|\beta|} A_{\alpha+\beta} A_{-\alpha} A_{-\beta}}{C_{\alpha, \beta}^{\alpha+\beta}} (h_\alpha, h_\beta)
\end{aligned}$$

Using these formulas, we can now see that each pair adds up to zero, and this completes the proof of the lemma. ■

This lemma shows that in order to study zero-weight super dynamical r-matrices with nonzero coupling constant, it suffices to solve the modified dynamical Yang-Baxter equation. This modified version of the dynamical Yang-Baxter equation:

$$Alt_s(ds) + [[s, s]] + \frac{\epsilon^2}{4} [[\Omega, \Omega]] = 0, \quad (3)$$

is an equation in $\mathfrak{g} \otimes \mathfrak{g} \otimes \mathfrak{g}$. The non-Casimir components are easily seen to be skew-symmetric with respect to signed permutations of factors, following arguments from the proof of Theorem 2.1. The Casimir part is also skew-symmetric with respect to signed permutations of factors, as can be seen by a straight-forward computation. Because of this symmetry, it is sufficient to look only at the $\mathfrak{h} \otimes \mathfrak{h} \otimes \mathfrak{h}$, $\mathfrak{h} \otimes \mathfrak{g}_\alpha \otimes \mathfrak{g}_{-\alpha}$, and $\mathfrak{g}_\alpha \otimes \mathfrak{g}_\beta \otimes \mathfrak{g}_{-\alpha-\beta}$ components in order to solve Equation 3.

The $\mathfrak{h} \otimes \mathfrak{h} \otimes \mathfrak{h}$ part comes only from the dynamical part (in other words, from $Alt_s(ds)$) and yields:

$$\frac{\partial D_{jk}}{\partial x_i} + \frac{\partial D_{ki}}{\partial x_j} + \frac{\partial D_{ij}}{\partial x_k} = 0,$$

which implies that $D = \sum_{i < j} D_{ij} dx_i \wedge dx_j$ is a closed differential 2-form.

The $\mathfrak{h} \otimes \mathfrak{g}_\alpha \otimes \mathfrak{g}_{-\alpha}$ part consists of two components. The part that comes from the non-Casimir part is the same as the whole of the $\mathfrak{h} \otimes \mathfrak{g}_\alpha \otimes \mathfrak{g}_{-\alpha}$ part

of the dynamical Yang-Baxter equation for a zero-weight super dynamical r-matrix with zero coupling constant, (see Section 2.3):

$$\begin{aligned}
 & Alt_s(ds) \big|_{\mathfrak{h} \otimes \mathfrak{g}_\alpha \otimes \mathfrak{g}_{-\alpha}} + \sum_{\alpha, \beta} (-1)^{|\alpha||\beta|} \delta_{\alpha, -\beta} \varphi_\alpha \varphi_\beta [e_\alpha, e_\beta] \otimes e_{-\alpha} \otimes e_{-\beta} \\
 &= \sum_{i, \alpha} \frac{\partial \varphi_\alpha}{\partial x_i} x_i \otimes e_\alpha \otimes e_{-\alpha} + \sum_{\alpha} (-1)^{|\alpha|} \varphi_\alpha \varphi_{-\alpha} (-A_\alpha) h_\alpha \otimes e_\alpha \otimes e_{-\alpha} \\
 &= \sum_{i, \alpha} \frac{\partial \varphi_\alpha}{\partial x_i} x_i \otimes e_\alpha \otimes e_{-\alpha} + \sum_{\alpha} A_\alpha \varphi_\alpha \varphi_\alpha h_\alpha \otimes e_\alpha \otimes e_{-\alpha}
 \end{aligned}$$

To this we need to add the Casimir component, i.e. the terms that come from $\frac{\epsilon^2}{4} [[\Omega, \Omega]]$:

$$\begin{aligned}
 [[\Omega, \Omega]] \big|_{\mathfrak{h} \otimes \mathfrak{g}_\alpha \otimes \mathfrak{g}_{-\alpha}} &= \sum_{\alpha, \beta} (-1)^{|\alpha||\beta|} \delta_{\alpha, -\beta} A_\alpha A_\beta [e_\alpha, e_\beta] \otimes e_{-\alpha} \otimes e_{-\beta} \\
 &= \sum_{\alpha} (-1)^{|\alpha|} A_\alpha A_{-\alpha} [e_\alpha, e_{-\alpha}] \otimes e_{-\alpha} \otimes e_\alpha \\
 &= \sum_{\alpha} (-1)^{|\alpha|} A_\alpha A_{-\alpha} A_{-\alpha} h_\alpha \otimes e_{-\alpha} \otimes e_\alpha \\
 &= \sum_{\alpha} A_{-\alpha} h_\alpha \otimes e_{-\alpha} \otimes e_\alpha \\
 &= - \sum_{\alpha} A_\alpha h_\alpha \otimes e_\alpha \otimes e_{-\alpha}
 \end{aligned}$$

and so the $\mathfrak{h} \otimes \mathfrak{g}_\alpha \otimes \mathfrak{g}_{-\alpha}$ part of Equation 3 is:

$$\sum_{i, \alpha} \frac{\partial \varphi_\alpha}{\partial x_i} x_i \otimes e_\alpha \otimes e_{-\alpha} + \sum_{\alpha} A_\alpha \varphi_\alpha^2 h_\alpha \otimes e_\alpha \otimes e_{-\alpha} - \frac{\epsilon^2}{4} \sum_{\alpha} A_\alpha h_\alpha \otimes e_\alpha \otimes e_{-\alpha}$$

which we can rewrite as:

$$\sum_{\alpha \in \Delta} \left(\sum_i \frac{\partial \varphi_\alpha}{\partial x_i} x_i + A_\alpha (\varphi_\alpha^2 - \frac{\epsilon^2}{4}) h_\alpha \right) \otimes e_\alpha \otimes e_{-\alpha}$$

For this term to vanish we must have, for all $\alpha \in \Delta$:

$$\sum_i \frac{\partial \varphi_\alpha}{\partial x_i} x_i + A_\alpha (\varphi_\alpha^2 - \frac{\epsilon^2}{4}) h_\alpha = 0.$$

We can rewrite this as:

$$d\varphi_\alpha + A_\alpha (\varphi_\alpha^2 - \frac{\epsilon^2}{4}) dh_\alpha = 0.$$

If we define μ_α by:

$$\mu_\alpha = \begin{cases} \sqrt{-1} & \text{if } \alpha \text{ is an odd positive root} \\ 1 & \text{otherwise} \end{cases}$$

then $\mu_\alpha^2 = A_\alpha$, and the equation we need to solve is:

$$d\varphi_\alpha + \mu_\alpha^2(\varphi_\alpha^2 - \frac{\epsilon^2}{4})dh_\alpha = 0.$$

We assume $\varphi_\alpha^2 \neq \frac{\epsilon^2}{4}$ and let $u_\alpha = \mu_\alpha \varphi_\alpha$. Also let $\epsilon_\alpha = \frac{\mu_\alpha \epsilon}{2}$. Separating variables to integrate we obtain:

$$\frac{1}{\mu_\alpha} \int \frac{du_\alpha}{\epsilon_\alpha^2 - u_\alpha^2} = \int dh_\alpha \Rightarrow u_\alpha = \mu_\alpha \varphi_\alpha = \epsilon_\alpha \coth(\epsilon_\alpha \mu_\alpha h_\alpha + C)$$

and we get:

$$\varphi_\alpha = \frac{\epsilon}{2} \coth\left(\frac{A_\alpha \epsilon}{2}(h_\alpha - \nu_\alpha)\right)$$

for some $\nu_\alpha \in \mathbb{C}$. Here h_α is viewed as a linear function on $\mathfrak{h}^*$ via $h_\alpha(\lambda) = (\alpha, \lambda)$.

Finally we look at the $\mathfrak{g}_\alpha \otimes \mathfrak{g}_\beta \otimes \mathfrak{g}_{-\alpha-\beta}$ part of Equation 3. There is no contribution from the dynamical part. The terms from $[[s, s]]$ are (see Section 2.3 for more details):

$$\begin{aligned} \sum_{\alpha, \beta} (-1)^{|\alpha||\beta|} \varphi_\alpha \varphi_\beta [e_\alpha, e_\beta] \otimes e_{-\alpha} \otimes e_{-\beta} + \sum_{\alpha, \beta} \varphi_\alpha \varphi_\beta e_\alpha \otimes [e_{-\alpha}, e_\beta] \otimes e_{-\beta} \\ + \sum_{\alpha, \beta} (-1)^{|\alpha||\beta|} \varphi_\alpha \varphi_\beta e_\alpha \otimes e_\beta \otimes [e_{-\alpha}, e_{-\beta}] \end{aligned}$$

and the coefficient for the $e_\alpha \otimes e_\beta \otimes e_{-\alpha-\beta}$ term coming from this component is:

$$(-1)^{|\beta||\alpha+\beta|} \varphi_{-\beta} \varphi_{\alpha+\beta} C_{-\beta, \alpha+\beta}^\alpha + \varphi_\alpha \varphi_{\alpha+\beta} C_{-\alpha, \alpha+\beta}^\beta + (-1)^{|\alpha||\beta|} \varphi_\alpha \varphi_\beta C_{-\alpha, -\beta}^{-\alpha-\beta}$$

We can simplify this further:

$$\begin{aligned} & (-1)^{|\beta||\alpha+\beta|} \varphi_{-\beta} \varphi_{\alpha+\beta} C_{-\beta, \alpha+\beta}^\alpha + \varphi_\alpha \varphi_{\alpha+\beta} C_{-\alpha, \alpha+\beta}^\beta + (-1)^{|\alpha||\beta|} \varphi_\alpha \varphi_\beta C_{-\alpha, -\beta}^{-\alpha-\beta} \\ &= (-1)^{|\beta||\alpha+\beta|} \varphi_{-\beta} \varphi_{\alpha+\beta} \frac{(-1)^{|\alpha||\beta|} (-1)^{|\beta|} A_{-\beta}}{C_{\alpha, \beta}^{\alpha+\beta}} (h_\alpha, h_\beta) \\ &+ \varphi_\alpha \varphi_{\alpha+\beta} \frac{-(-1)^{|\alpha|} A_{-\alpha}}{C_{\alpha, \beta}^{\alpha+\beta}} (h_\alpha, h_\beta) \\ &+ (-1)^{|\alpha||\beta|} \varphi_\alpha \varphi_\beta \frac{(-1)^{|\alpha||\beta|} (-1)^{|\alpha|+|\beta|} A_{\alpha+\beta} A_{-\alpha} A_{-\beta}}{C_{\alpha, \beta}^{\alpha+\beta}} (h_\alpha, h_\beta) \\ &= -\varphi_\beta \varphi_{\alpha+\beta} \frac{A_\beta}{C_{\alpha, \beta}^{\alpha+\beta}} (h_\alpha, h_\beta) - \varphi_\alpha \varphi_{\alpha+\beta} \frac{A_\alpha}{C_{\alpha, \beta}^{\alpha+\beta}} (h_\alpha, h_\beta) \\ &+ \varphi_\alpha \varphi_\beta \frac{A_{\alpha+\beta} A_\alpha A_\beta}{C_{\alpha, \beta}^{\alpha+\beta}} (h_\alpha, h_\beta) \end{aligned}$$

The Casimir component is:

$$\begin{aligned} \frac{\epsilon^2}{4} \left(\sum_{\alpha, \beta} (-1)^{|\alpha||\beta|} A_\alpha A_\beta [e_\alpha, e_\beta] \otimes e_{-\alpha} \otimes e_{-\beta} + \sum_{\alpha, \beta} A_\alpha A_\beta e_\alpha \otimes [e_{-\alpha}, e_\beta] \otimes e_{-\beta} \right. \\ \left. + \sum_{\alpha, \beta} (-1)^{|\alpha||\beta|} A_\alpha A_\beta e_\alpha \otimes e_\beta \otimes [e_{-\alpha}, e_{-\beta}] \right) \end{aligned}$$

and the coefficient in front of the $e_\alpha \otimes e_\beta \otimes e_{-\alpha-\beta}$ term coming from this component is:

$$\begin{aligned} \frac{\epsilon^2}{4} \left((-1)^{|\beta||\alpha+\beta|} A_{-\beta} A_{\alpha+\beta} C_{-\beta, \alpha+\beta}^\alpha + A_\alpha A_{\alpha+\beta} C_{-\alpha, \alpha+\beta}^\beta \right. \\ \left. + (-1)^{|\alpha||\beta|} A_\alpha A_\beta C_{-\alpha, -\beta}^{-\alpha-\beta} \right) \end{aligned}$$

which we can simplify as follows:

$$\begin{aligned} & \frac{\epsilon^2}{4} \left((-1)^{|\beta||\alpha+\beta|} A_{-\beta} A_{\alpha+\beta} C_{-\beta, \alpha+\beta}^\alpha + A_\alpha A_{\alpha+\beta} C_{-\alpha, \alpha+\beta}^\beta \right. \\ & \quad \left. + (-1)^{|\alpha||\beta|} A_\alpha A_\beta C_{-\alpha, -\beta}^{-\alpha-\beta} \right) \\ &= \frac{\epsilon^2}{4} \left((-1)^{|\alpha||\beta|} A_\beta A_{\alpha+\beta} C_{-\beta, \alpha+\beta}^\alpha + A_\alpha A_{\alpha+\beta} C_{-\alpha, \alpha+\beta}^\beta \right. \\ & \quad \left. + (-1)^{|\alpha||\beta|} A_\alpha A_\beta C_{-\alpha, -\beta}^{-\alpha-\beta} \right) \\ &= \frac{\epsilon^2}{4} \left(((-1)^{|\alpha||\beta|} A_\beta A_{\alpha+\beta} \frac{(-1)^{|\alpha||\beta|} (-1)^{|\beta|} A_{-\beta}}{C_{\alpha, \beta}^{\alpha+\beta}} (h_\alpha, h_\beta) \right. \\ & \quad + A_\alpha A_{\alpha+\beta} \frac{-(-1)^{|\alpha|} A_{-\alpha}}{C_{\alpha, \beta}^{\alpha+\beta}} (h_\alpha, h_\beta) \\ & \quad \left. + (-1)^{|\alpha||\beta|} A_\alpha A_\beta \frac{(-1)^{|\alpha||\beta|} (-1)^{|\alpha|+|\beta|} A_{\alpha+\beta} A_{-\alpha} A_{-\beta}}{C_{\alpha, \beta}^{\alpha+\beta}} (h_\alpha, h_\beta) \right) \\ &= \frac{\epsilon^2}{4} \left(((-1)^{|\alpha||\beta|} A_{\alpha+\beta} \frac{(-1)^{|\alpha||\beta|}}{C_{\alpha, \beta}^{\alpha+\beta}} (h_\alpha, h_\beta) + A_{\alpha+\beta} \frac{-1}{C_{\alpha, \beta}^{\alpha+\beta}} (h_\alpha, h_\beta) \right. \\ & \quad \left. + (-1)^{|\alpha||\beta|} \frac{(-1)^{|\alpha||\beta|} A_{\alpha+\beta}}{C_{\alpha, \beta}^{\alpha+\beta}} (h_\alpha, h_\beta) \right) \\ &= \frac{\epsilon^2}{4} \left(\frac{A_{\alpha+\beta}}{C_{\alpha, \beta}^{\alpha+\beta}} (h_\alpha, h_\beta) \right) \end{aligned}$$

Therefore the coefficient of the $e_\alpha \otimes e_\beta \otimes e_{-\alpha-\beta}$ term is:

$$\begin{aligned} & - \varphi_\beta \varphi_{\alpha+\beta} \frac{A_\beta}{C_{\alpha,\beta}^{\alpha+\beta}}(h_\alpha, h_\beta) - \varphi_\alpha \varphi_{\alpha+\beta} \frac{A_\alpha}{C_{\alpha,\beta}^{\alpha+\beta}}(h_\alpha, h_\beta) \\ & + \varphi_\alpha \varphi_\beta \frac{A_{\alpha+\beta} A_\alpha A_\beta}{C_{\alpha,\beta}^{\alpha+\beta}}(h_\alpha, h_\beta) + \frac{\epsilon^2}{4} \left(\frac{A_{\alpha+\beta}}{C_{\alpha,\beta}^{\alpha+\beta}}(h_\alpha, h_\beta) \right) \end{aligned}$$

This is equal to zero if and only if:

$$\varphi_\beta \varphi_{\alpha+\beta} A_\beta + \varphi_\alpha \varphi_{\alpha+\beta} A_\alpha = \varphi_\alpha \varphi_\beta A_{\alpha+\beta} A_\alpha A_\beta + \frac{\epsilon^2}{4} A_{\alpha+\beta},$$

or equivalently:

$$A_{\alpha+\beta} \varphi_\alpha \varphi_\beta + \frac{\epsilon^2}{4} A_{\alpha+\beta} A_\alpha A_\beta = \varphi_{\alpha+\beta} (A_\alpha \varphi_\beta + A_\beta \varphi_\alpha) \quad (4)$$

where we use $A_\alpha^2 = 1$.

Let $X = \{\alpha \in \Delta \mid \varphi_\alpha^2 \neq \frac{\epsilon^2}{4}\}$. Then clearly X is closed under changing signs because s satisfies the unitarity condition. Equation 4 implies that X is also closed under addition. If $\alpha, \beta \in X$ are two positive roots, then:

$$\begin{aligned} \varphi_\alpha &= \frac{\epsilon}{2} \coth \left(\frac{A_\alpha \epsilon}{2} (h_\alpha - \nu_\alpha) \right) \\ \varphi_\beta &= \frac{\epsilon}{2} \coth \left(\frac{A_\beta \epsilon}{2} (h_\beta - \nu_\beta) \right) \end{aligned}$$

and Equation 4 implies:

$$\varphi_{\alpha+\beta} = \frac{\epsilon}{2} A_{\alpha+\beta} \coth \left(\frac{\epsilon}{2} (h_\alpha - \nu_\alpha) + \frac{\epsilon}{2} (h_\beta - \nu_\beta) \right)$$

is not constant, so we must have:

$$\varphi_{\alpha+\beta} = \frac{\epsilon}{2} \coth \left(\frac{A_{\alpha+\beta} \epsilon}{2} (h_{\alpha+\beta} - \nu_{\alpha+\beta}) \right).$$

Since $\coth(-x) = -\coth(x)$, this implies also that:

$$\nu_{\alpha+\beta} = \nu_\alpha + \nu_\beta.$$

Using $\varphi_{-\alpha} = -(-1)^{|\alpha|} \varphi_\alpha$ we can also see that $\nu_{-\alpha} = -\nu_\alpha$. Therefore we can conclude that there is some $\nu \in \mathfrak{h}^*$ such that $\nu_\alpha = (\alpha, \nu)$ for all $\alpha \in X$.

Thus we have proved that r is a super dynamical r-matrix with zero weight and coupling constant ϵ if and only if $s = r - \frac{\epsilon}{2} \Omega$ is of the following form:

$$s(\lambda) = \sum_{i,j=1}^N D_{ij}(\lambda) x_i \otimes x_j + \sum_{\alpha \in \Delta} \varphi_\alpha e_\alpha \otimes e_{-\alpha}$$

and this finishes the proof of Theorem 2.2. ■

3. EXAMPLES OF SUPER DYNAMICAL r -MATRICES

Example 3.1. Let $\mathfrak{g}$ be a simple Lie superalgebra with nondegenerate Killing form. For any choice of a triangular decomposition of $\mathfrak{g}$, the constant r -matrix:

$$r = \frac{\epsilon}{2} \sum_i (x_i \otimes x_i^*) + \epsilon \sum_{\alpha \in \Delta^+} (e_{-\alpha} \otimes e_{\alpha})$$

and its super twist:

$$T_s(r) = \frac{\epsilon}{2} \sum_i (x_i \otimes x_i^*) + \epsilon \sum_{\alpha \in \Delta^+} \left((-1)^{|\alpha|} e_{\alpha} \otimes e_{-\alpha} \right)$$

are both zero-weight super dynamical r -matrices with coupling constant ϵ . They correspond to $X = \emptyset$ and $D \equiv 0$ in Theorem 2.2. We note that these r -matrices are in fact solutions to the classical Yang-Baxter equation. In fact, r can clearly be constructed by the main construction theorem of [8] with the trivial admissible triple $(\Gamma_1, \Gamma_2, \tau)$ where $\Gamma_i = \emptyset$ and $\tau = 0$.

Remark. The Cartan components of the r -matrices above involve $x_i^* \in \mathfrak{h}$, the elements of the dual basis of the basis $\{x_i\}$ for $\mathfrak{h}$, because it may not be always possible to choose an orthonormal basis for $\mathfrak{h}$ in the super case.

Remark. Clearly the only solution of the classical Yang-Baxter equation constructible by the main construction theorem of [8] that we can also construct by Theorem 2.2 is r given above. To reach any other r -matrix constructible by this method, i.e. one that involves a nontrivial admissible triple, we would need to drop the zero weight condition.

Example 3.2. For a non-constant dynamical example, consider the super dynamical r -matrix constructed by Theorem 2.2 using $X = \Delta$, $D \equiv 0$ and $\nu = 0$:

$$r(\lambda) = \frac{\epsilon}{2} \Omega + \sum_{\alpha \in \Delta} \frac{\epsilon}{2} \coth \left(\frac{\epsilon}{2} (\alpha, \lambda) \right) (-1)^{|\alpha|} (e_{\alpha}, e_{-\alpha}) (e_{\alpha} \otimes e_{-\alpha})$$

For a given triangular decomposition this can be rewritten as:

$$r(\lambda) = \frac{\epsilon}{2} \Omega + \sum_{\alpha \in \Delta^+} \frac{\epsilon}{2} \coth \left(\frac{\epsilon}{2} (\alpha, \lambda) \right) \left((-1)^{|\alpha|} e_{\alpha} \otimes e_{-\alpha} - e_{-\alpha} \otimes e_{\alpha} \right).$$

For this triangular decomposition, if we take the limit as $\lambda \rightarrow \infty$ while always staying positive, then we see that:

$$r(\lambda) \longrightarrow \frac{\epsilon}{2} \Omega + \sum_{\alpha \in \Delta^+} \frac{\epsilon}{2} \left((-1)^{|\alpha|} e_{\alpha} \otimes e_{-\alpha} - e_{-\alpha} \otimes e_{\alpha} \right)$$

which is exactly the r -matrix $T_s(r)$ of the previous example. If we take a similar limit as $|\lambda| \rightarrow \infty$ while staying always negative, we see that:

$$r(\lambda) \longrightarrow \frac{\epsilon}{2} \Omega + \sum_{\alpha \in \Delta^+} \frac{\epsilon}{2} \left(e_{-\alpha} \otimes e_{\alpha} - (-1)^{|\alpha|} e_{\alpha} \otimes e_{-\alpha} \right)$$

which is exactly the r -matrix r of the previous example.

Hence we can conclude that this particular super dynamical r -matrix extrapolates different solutions of the classical Yang-Baxter equation associated to the trivial admissible triple, labeled by different triangular decompositions.

4. CONCLUSION

The two theorems proven in this paper are the super versions of Theorem 3.2 and Theorem 3.10 in [6]. Their proofs were very clearly inspired by the non-graded case. A careful eye will discern the difficulties that are peculiar to the super case in the proofs provided here.

For a classification of all super dynamical r -matrices, one needs further study. Schiffmann's results in [10] provide a good template. It is this author's belief that the constructive part of Schiffmann's results will carry over to the super case with certain modifications. In light of recent results, one expects to see some divergence from the non-graded theory when it comes to the classification part; in fact the example studied in [9] will be one of the challenges to a direct generalization.

REFERENCES

- [1] Belavin, A. A., Drinfeld, V. G.; "*Solutions of the Classical Yang-Baxter Equation and Simple Lie Algebras*", *Funct. Anal. Appl.* **16** (1982), pp.159–180.
- [2] Belavin, A. A., Drinfeld, V. G.; "*Triangle Equation and Simple Lie Algebras*", *Soviet Scientific Reviews Sect. C* **4** (1984), pp.93–165.
- [3] Etingof, P.; "*On the Dynamical Yang-Baxter Equation*", *Proceedings of the International Congress of Mathematicians, Vol. II (Beijing, 2002)*, Higher Ed. Press, 2002, pp.555–570.
- [4] Etingof, P., Schedler, T., Schiffmann, O.; "*Explicit quantization of dynamical r -matrices for finite dimensional semisimple Lie algebras*"; *J. Amer. Math. Soc.* **13** (2000), no. 3, pp.595–609.
- [5] Etingof, P., Schiffmann, O.; *Lectures on Quantum Groups*, International Press, 1998.
- [6] Etingof, P., Varchenko, A.; "*Geometry and Classification of Solutions of the Classical Dynamical Yang-Baxter Equation*", *Comm. Math. Phys.* **192** (1998), no. 1, pp.77–120.
- [7] Geer, N.; "*Etingof-Kazhdan Quantization of Lie Superalgebras*"; e-arXiv preprint, [arXiv:math.QA/0409563](https://arxiv.org/abs/math.QA/0409563)
- [8] Karaali, G.; "*Constructing r -matrices on Simple Lie Superalgebras*", *J. Algebra* **282** (2004), no.1, pp.83–102.
- [9] Karaali, G.; "*A New Lie Bialgebra Structure on $sl(2,1)$* ", submitted.
- [10] Schiffmann, O.; "*On Classification of Dynamical r -matrices*", *Math. Res. Lett.* **5** (1998), pp.13–30.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, SANTA BARBARA, CA 93106

E-mail address: gizem@math.ucsb.edu